

Centre de Physique Théorique* - CNRS - Luminy, Case 907
F-13288 Marseille Cedex 9 - France

THE CONVERSION OF GRAVITONS INTO PHOTONS IN TE_{mo} MODE

Nguyen Suan Han¹, Hoang Ngoc Long² and Dang Van Soa¹

Abstract

We consider the conversion of gravitons into photons in the TE_{mo} mode. Cross sections in different directions are given.

November 1994
CPT-94/P.3080

anonymous ftp or gopher: cpt.univ-mrs.fr

*Unité Propre de Recherche 7061

¹Permanent address: Department of Theoretical Physics, Hanoi State University,
P. O. Box 600, BoHo, Hanoi, 10000, Vietnam

²Institute of Theoretical Physics, P. O. Box 429, BoHo, Hanoi10000, Vietnam.

1 Introduction

The study on the interaction between electromagnetic (EM) and gravitational fields is a significant part of the research on gravitational radiation [1 – 3]. Based on the present technical level in the laboratory, it has been proved by means of mechanics that it is as yet difficult to generate gravitational wave (GW) which are strong enough to be detected by current detectors. Therefore physicists have transferred their interest to strong EM field and taken it as one of the possible sources in the laboratory.

In previous paper [4] we have considered the conversion of gravitons into photons in the periodic external EM field, namely in the TE_{10} mode. In this paper we consider the conversion of gravitons into photons in the TE_{mo} mode [5] :

2 Cross sections

We consider the case of an alternating EM field in a rectangular cavity with inner dimensions a , b , propagating in $z = -c/2$ and $z = c/2$. To simplify we use a system of units in which $\hbar = c = 1$, choose the nontrivial solution TE_{mo} mode [5]:

$$\begin{aligned} H_z &= H_0 \cos\left(\frac{m\pi x}{a}\right) e^{ikz-i\omega t} , \\ H_x &= -\frac{ika}{m\pi} H_0 \sin\left(\frac{m\pi x}{a}\right) e^{ikz-i\omega t} , \\ E_y &= i\frac{\omega a}{m\pi} H_0 \sin\left(\frac{m\pi x}{a}\right) e^{ikz-i\omega t} , \end{aligned} \quad (1)$$

where $k = w/v$.

Let us consider the conversion of the graviton g with momentum p into a photon γ with momentum q in the above mentioned external EM field. For the above mentioned process we get the matrix element:

$$\begin{aligned} \langle q|M_{g\gamma}|p \rangle &= \frac{k}{4(2\pi)^2\sqrt{q_0p_0}} \epsilon^i(\vec{q}, \sigma) \left[B\epsilon^{j2}(\vec{p}, \tau)\eta_{ij}q_0 + (\eta_{i2}q_j - \eta_{ij}q_y)(D\epsilon^{j3}(\vec{p}, \tau) - C\epsilon^{j1}(\vec{p}, \tau)) \right. \\ &\quad \left. + (C\eta_{i1} - D\eta_{i3})\epsilon^{j2}(\vec{p}, \tau)q_j + \epsilon^{j2}(\vec{p}, \tau)\eta_{ij}(Dq_z - Cq_x) \right] , \end{aligned} \quad (2)$$

where $k = \sqrt{16\pi G}$, $q_o = p_o + \omega$ and

$$\begin{aligned} B &= \frac{(-1)^{m-1}/28a\omega H_0(p_x - q_x)}{m\pi[(p_x - q_x)^2 - \frac{m^2\pi^2}{a^2}(p_y - q_y)(p_z - q_z + k)]} \cos\frac{a}{2}(p_x - q_x) \sin\frac{b}{2}(p_y - q_y) \sin\frac{c}{2}(p_z - q_z + k), \\ C &= -\frac{(-1)^{\frac{m-1}{2}}8m\pi H_0}{a[(p_x - q_x)^2 - \frac{m^2\pi^2}{a^2}(p_y - q_y)(p_z - q_z + k)]} \cos\frac{a}{2}(p_x - q_x) \sin\frac{b}{2}(p_y - q_y) \sin\frac{c}{2}(p_z - q_z + k), \\ D &= -\frac{(-1)^{(m-1)/2}8kaH_0(p_x - q_x)}{m\pi[(p_x - q_x)^2 - \frac{m^2\pi^2}{a^2}(p_y - q_y)(p_z - q_z + k)]} \cos\frac{a}{2} \sin\frac{b}{2}(p_y - q_y) \sin\frac{c}{2}(p_z - q_z + k) \end{aligned}$$

for $m = 2k - 1, k = 1, 2, 3, \dots$ and

$$B = \frac{-(-1)^{m/2} 8\omega H_0}{[(p_x - q_x)^2 - \frac{m^2 \pi^2}{a^2}](p_y - q_y)(p_z - q_z + k)} \sin \frac{a}{2}(p_x - q_x) \sin \frac{b}{2}(p_y - q_y) \sin \frac{c}{2}(p_z - q_z + k),$$

$$C = \frac{(-1)^{m/2} 8H_0(p_x - q_x)}{[(p_x - q_x)^2 - \frac{m^2 \pi^2}{a^2}](p_y - q_y)(p_z - q_z + k)} \sin \frac{a}{2}(p_x - q_x) \sin \frac{b}{2}(p_y - q_y) \sin \frac{c}{2}(p_z - q_z + k),$$

$$D = \frac{(-1)^{m/2} 8kH_0}{[(p_x - q_x)^2 - \frac{m^2 \pi^2}{a^2}](p_y - q_y)(p_z - q_z + k)} \sin \frac{a}{2}(p_x - q_x) \sin \frac{b}{2}(p_y - q_y) \sin \frac{c}{2}(p_z - q_z + k).$$

for $m = 2k$.

In the following, we assume that $v = 1$ and $\omega \ll p$. We also retain the terms to the first order in ω in the result.

When the momentum of the graviton is parallel to the z axis (the direction of the external field propagation), the differential cross section, in the context of the previous statement, vanishes ($\sim O(\omega^2)$) when $\theta \approx 0$ and when $\theta = \pi/2$.

Now we consider the case in which the momentum of graviton is parallel to the x axis, i. e. $p^\mu = (p, p, 0, 0)$. We have

$$\begin{aligned} \frac{d\sigma(g \rightarrow \gamma)}{d\Omega'} &= \frac{k^2 q}{64(2\pi)^2 p} \left\{ q^2 C^2 (1 + \cos^2 \theta - 2 \sin^2 \theta \cos^2 \varphi') - \right. \\ &\quad 2C \left[q \cos \theta (1 + \cos^2 \theta + \sin^2 \theta) (q_0 B + q D \sin \theta \sin \varphi') \right. \\ &\quad \left. \left. - q^2 \sin(2\theta) \sin \varphi' \right] \right\}. \end{aligned} \quad (3)$$

From (3) it follows that ($m = 2k - 1$)

$$\frac{d\sigma(g \rightarrow \gamma)}{d\Omega'} = \frac{k^2 H_0^2 V^2 p^2}{2m^2 (2\pi)^4} \left(1 + 3 \frac{\omega}{p} \right) \quad (4)$$

for $\theta \approx 0$, and

$$\frac{d\sigma(g \rightarrow \gamma)}{d\Omega'} = \frac{k^2 H_0^2 b^2 m^2}{16a^2 (p^2 - \frac{m^2 \pi^2}{a^2})^2} \left(1 + 3 \frac{\omega}{p} \right) \cos^2 \left(\frac{ap}{2} \right) 2 \sin^2 \left[\frac{c(p + 2k)}{2} \right] \quad (5)$$

for $\theta = \frac{\pi}{2}, \theta = \frac{\pi}{2}$ and for $m = 2k$

$$\frac{d\sigma(g \rightarrow \gamma)}{d\Omega'} = 0 \quad (6)$$

for the $\theta \approx 0$, and

$$\frac{d\sigma(g \rightarrow \gamma)}{d\Omega'} = \frac{k^2 H_0^2 b^2 p^2}{4(2\pi)^4 (p^2 - \frac{m^2 \pi^2}{a^2})} \left(1 + 3 \frac{\omega}{p} \right) \sin^2 \left(\frac{ap}{2} \right) 2 \sin^2 \left[\frac{c(p + 2k)}{2} \right] \quad (7)$$

for $\theta = \pi/2, \varphi = \pi/2$.

Next, we consider the case in which the momentum of the graviton is parallel to the y axis, i. e. $p^\mu = (p, 0, p, 0)$. We then have

$$\frac{d\sigma(g \rightarrow \gamma)}{d\Omega''} = \frac{k^2 q^3}{64(2\pi)^2 p} (C^2 + D^2) (3 \cos^2 \theta - 1) . \quad (8)$$

From (8) it follows that the differential cross section vanishes in the region $120^\circ \gg \theta \gg 60^\circ$ and gets the largest values.

$$\frac{d\omega(g \rightarrow \gamma)}{d\Omega''} = \frac{k^2 H_c^2 V^2 p^2}{2(2\pi)^4 m^2} \left(1 + 3 \frac{\omega}{p} \right) \quad (9)$$

for $\theta \approx 0$ and $m = 2k - 1$.

3 Acknowledgements

We are grateful to Profs. B.M. Barbashov, V.N. Pervushin, A.M. Khvedelidze, Dao Vong Duc, Tran Huu Phat for valuable discussions. One of us (NSH) is indebted to Prof. P. Chiappetta for financial support during his stay at the Centre de Physique Théorique in Marseille and warm hospitality.

References

- [1] I. M. Pinto and G. Rotoli, in "Proceedings of the 8th Italian Conference on General Relativity and Gravitational Physics", eds. M. G. Cerdonio, R. Cianci, M. Francaviglia and M. Tolli. World Scientific, 1989, p. 560. E. Amaldi and F. Fuligni, Report and contribution papers to workshop C1, in "General Relativity and Gravitation", eds. B. Bertotti F. DE Felice, and A. Pascolini, D. Reidel, 1984. L. P. Grishchuk, in "Proceedings of the 9th Inter. Conf. on General Relativity and Gravitation", Jena 1980, ed. E. Schnutser, Cambridge Univ. Press. 255(1983) and references therein.
- [2] D. Boccaletti, V. De Sabbata, P. Fortini, and C. Gualdi, Nuovo Cimento 70B, (1970)120. D. Boccaletti and F. Occhionero, Lett. Nuovo Cimento, 2(1971)549. J. Weber, General Relativity and Gravitational Waves, Interscience, N. Y. (1961), N. V. Mitskievich, Physical Fields in General Relativity, Nauka, Moscow, (196). L. P. Grishchuk and A. G. Polnarev, in " General Relativity and Gravitation", vol. 2, ed. A. Held. Plenum Publishing Corporation, New York, 1980, p. 393. G. Papini and S. R. Valluri, Phys. Rep. 33(1978)51, L. P. Grishchuk and M. V. Sazhin, Sov. Phys. , JETP, 41(1975)787, V. B. Braginsky et al. , Ibit. 38(1973)865,
- [3] H. N. Long, L. K. Huong. Mod. Phys. Lett. A6(1991)2315,
- [4] H. N. Long, D. V. Soa, T. A. Tuan, Phys. Lett. A, vol. 186, (1994) 382,
- [5] J. D. Jackson, Classical Electrodynamics, John Wiley & Sons, Inc. , 1975, Sect. 8. 4, American Institute of Physics Handbook, ed. D. E. Gray, 3rd Edition, McGraw Hil, New York (1972)p. 5.